

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

Journal of Oral Biosciences

journal homepage: www.elsevier.com/locate/job

Original Article

Microbial composition of dental plaque microflora on first molars with orthodontic bands and brackets, and the acidogenic potential of these bacteria

Ryo Komori^a, Takuichi Sato^{b,*}, Teruko Takano-Yamamoto^a, Nobuhiro Takahashi^b^a Division of Orthodontics and Dentofacial Orthopedics, Tohoku University Graduate School of Dentistry, 4-1 Seiryō-machi, Aoba-ku, Sendai 980-8575, Japan^b Division of Oral Ecology and Biochemistry, Tohoku University Graduate School of Dentistry, 4-1 Seiryō-machi, Aoba-ku, Sendai 980-8575, Japan

ARTICLE INFO

Article history:

Received 22 November 2011

Received in revised form

4 January 2012

Accepted 23 January 2012

Available online 23 May 2012

Keywords:

16S ribosomal RNA

Acidogenic potential

Orthodontic bands

Orthodontic brackets

Dental plaque

ABSTRACT

Purpose: In order to elucidate the characteristics of orthodontic appliance-associated dental plaque, this study aimed to profile the microflora and to estimate the acidogenic potential of supragingival plaque on first molars with orthodontic bands or brackets or without appliances.

Material and methods: Supragingival plaque samples from the surface of upper and lower first molars with orthodontic bands or brackets or without appliances in 6 subjects (age, 11–30 years) were cultured anaerobically on blood agar plates. Isolated bacteria were identified by 16S rRNA sequencing. The acidogenicity of isolated bacteria was examined using fastidious anaerobe agar plates containing bromocresol.

Results: Bacterial growth (log CFU/mg) was 6.6 ± 6.5 , 6.9 ± 7.1 , and 7.4 ± 7.6 in samples obtained from molars with bands, brackets, and without appliances, respectively. *Actinomyces* (43.5 and 40.0%) and *Streptococcus* (23.5 and 34.7%) were the predominant species present on molars with orthodontic brackets and without appliances, respectively. In contrast, the proportion of *Streptococcus* (44.4%) was higher than that of *Actinomyces* (17.6%) on molars with orthodontic bands ($P < 0.01$). The proportions of acidogenic bacteria in plaque samples from molars with bands, brackets, and without appliances were 74.5%, 71.3%, and 81.6%, respectively, although these differences were not statistically significant.

Discussion and conclusion: These results indicate that there are differences in the microbial composition and acidogenic potential of supragingival plaque from first molars with bands, brackets, or without appliances, and suggest that supragingival plaque on teeth with brackets carries predominantly periodontitis-associated bacteria but less caries-associated bacteria.

© 2012 Japanese Association for Oral Biology. Published by Elsevier B.V. All rights reserved.

1. Introduction

More than 600 bacterial species comprise the plaque microflora that exists on surfaces within the oral cavity^{1,2}. Numerous environmental changes within the oral cavity can lead to changes in the composition of plaque microflora³.

The risk of oral diseases, including dental caries, gingivitis, and periodontitis, increases when various orthodontic appliances are attached to the enamel surface of teeth⁴. For instance, the risk of dental caries is reported to increase as the proportion of caries-related bacteria increases in the plaque microflora⁵. Caries-related bacteria, including *Streptococcus mutans* and *Lactobacillus*^{6–10}, and periodontitis-associated bacteria, such as *Eubacterium*, *Fusobacterium*, and *Treponema*^{10–16}, have been detected in orthodontic

appliance-associated dental plaque by PCR and other molecular techniques. However, with the use of such molecular techniques alone, it is not only difficult to clarify the microbial composition of plaque microflora but also to determine the amount of live bacteria within the microflora. To date, no comprehensive studies have been carried out where anaerobic culture is used in combination with 16S rRNA gene sequencing for the accurate comparison of bacteria in plaque microflora associated with orthodontic appliances. Furthermore, despite the fact that plaque biofilm is considered to carry an increased risk of cariogenicity^{17–19}, we were unable to find any previous studies that have used bacterial microflora isolates to determine its acid productivity in patients with orthodontic attachments.

Therefore, the aim of this study was to determine the bacterial levels, composition, and acidogenic potential of plaque microflora on upper and lower first molars without appliances or with bands or brackets, in order to elucidate the characteristics of orthodontic appliance-associated dental plaque.

* Corresponding author. Tel.: +81 22 717 8295; fax: +81 22 717 8297.

E-mail address: tak@m.tohoku.ac.jp (T. Sato).

2. Materials and Methods

2.1. Subjects

Six periodontally healthy subjects (4 females and 2 males; aged 11–30 y; mean, 21.5 y) undergoing orthodontic treatment at the Clinical Division of Orthodontics, Tohoku University Hospital, Sendai, Japan, were randomly selected for this study. Subjects were medically healthy, had received no antibiotics for 3 weeks before sampling, and were not taking immunosuppressive medication, anti-inflammatory drugs, or radiation therapy. Furthermore, they had no anamnesis of pregnancy, genetic diseases, or smoking habits. There were no instances of gingival sulcus depth greater than 3 mm or alveolar bone loss. Informed consent was obtained from all subjects, and the study was approved by the Ethics Review Board of Tohoku University, Graduate School of Dentistry, Sendai, Japan.

2.2. Orthodontic bands and brackets

Orthodontic bands (Tomy, Tokyo, Japan) made of stainless steel (sterilized) were fixed on the first molars using self-curing, glass ionomer cement (Fuji IX, GC, Tokyo, Japan). A dental probe and scaler were used for cleaning residual cement from the bands. Orthodontic brackets (Dentsply-Sankin, Tokyo, Japan) made of stainless steel (sterilized) were fixed on the surface of first molars using self-curing, adhesive resin (Super Bond, Sun Medical, Moriyama, Japan). The average sampling periods after fixation of orthodontic bands and brackets were 3.8 months (range, 2.6–5.0 months) and 3.1 months (range, 0.5–9.3 months), respectively (Table 1). Until the time of sampling, subjects were recalled once a month and were given oral hygiene instructions in order to ensure adequate tooth brushing technique. Subjects did not use mouth rinses (e.g., chlorhexidine) during the trial period, but there were no restrictions with regard to diet. Subjects were asked to refrain from tooth brushing on the morning before plaque sampling.

2.3. Sampling procedure

After sampling sites were isolated with cotton rolls, supragingival plaque samples (buccal sites) on the upper or lower first molars with orthodontic bands or brackets (on the tooth surface between the area above gingival margin and below the bands or brackets) were collected with sterilized toothpicks. Control samples from first molars without orthodontic bands or brackets were obtained at the same time as the test samples in subjects 1, 5, and 6. In addition, control samples were obtained from subject 3 one month before fixation of orthodontic bands and brackets, and from subjects 2 and 4 one month after the removal of orthodontic brackets.

2.4. Isolation of bacteria

Samples were transported in tightly screw-capped tubes, weighed, and transferred as soon as possible to an anaerobic glove box (Model AZ-Hard, Hirasawa, Tokyo, Japan) containing 80% N₂, 10% H₂, and 10% CO₂. Each sample was then suspended at a concentration of 1.0 mg/mL in sterilized 40 mM potassium phosphate buffer (pH 7.0) and was dispersed with a Teflon homogenizer. Serial 10-fold dilutions (0.1 mL each) were spread onto the surface of CDC anaerobic 5% sheep blood agar (BD, Franklin Lakes, NJ) plates, and were incubated in the anaerobic glove box at 37 °C for 7 d. After incubation, colony-forming units (CFU) on the plates were enumerated, and all colonies from plates having fewer than 100 colonies (mean, 16.5; range, 4–24

Table 1
Clinical features of subjects and bacterial growth in samples.

	Orthodontic bands						Orthodontic brackets						Without appliances						
	1	3	5	6	6		Mean ± SD	2	3	4		Mean ± SD	1	2	3	4	5	6	Mean ± SD
Subjects	26	20	11	30	30	female	21.8 ± 8.3	25	20	17	Female	20.7 ± 4.0	26	25	20	17	11	30	19.5 ± 6.9
Age (years)	Female	Male	Male	UL	UL	female		Female	Male	Female		Female	Female	Female	Male	Female	Male	Female	
Gender	UR	UL	UL	UL	UL	UL		UR	UL	UL	UR	UR	UR	UL	UR	UL	LR	LR	
Sampled tooth (first molar) ^a	5.0	5.0	2.6	3.0	4.0	4.0	3.8 ± 1.0	0.7	0.7	0.5	0.7	3.1 ± 3.8	I ^c	II ^c	III ^c	II ^c	I ^c	I ^c	
Sampling period ^b	6.7	6.7	6.3	6.3	7.0	6.7	6.2	6.6 ± 6.5	6.5	6.5	6.5	6.8	6.9 ± 7.1	6.5	5.9	6.5	7.7	6.8	7.3
Log (CFU/mg)																			7.4 ± 7.6

^a UR, upper right; UL, upper left; LR, lower right; LL, lower left.

^b Sampling period (months) after attachment of orthodontic appliances.

^c I, concurrently; II, 1 month after removal; III, 1 month before application.

colonies) were sub-cultured onto CDC plates. All plates, media, buffer solutions, and experimental instruments were kept in the anaerobic glove box for at least 24 h before use. To ensure strictly anaerobic conditions in the glove box, the reduction of methyl viologen (-446 mV) was carefully monitored whenever experimental procedures were carried out.

2.5. DNA extraction and identification of bacteria by 16S rRNA gene sequencing

Genomic DNA was extracted from each single colony using InstaGene Matrix kits (BioRad Laboratories, Richmond, CA), in accordance with the manufacturer's instructions.

The 16S rRNA gene sequences were amplified by PCR using universal primers 27 F and 1492 R^{20,21} and Taq DNA polymerase (HotStar Taq Master Mix, Qiagen GmbH, Hilden, Germany), according to the manufacturer's instructions. Primer sequences were as follows: 27 F, 5'-AGA GTT TGA TCM TGG CTC AG-3' and 1492 R, 5'-TAC GGY TAC CTT GTT ACG ACT T-3'. Amplification proceeded using a PCR Thermal Cycler MP (Takara Biomedicals, Ohtsu, Japan) programmed as follows: 15 min at 95 °C for initial heat activation; 30 cycles of 1 min at 94 °C for denaturation, 1 min at 55 °C for annealing, and 1.5 min at 72 °C for extension; and 10 min at 72 °C for final extension. PCR products were separated on 1% agarose gels (High Strength Analytical Grade Agarose, BioRad Laboratories) in Tris-borate EDTA buffer (100 mM Tris, 90 mM borate, 1 mM EDTA; pH 8.4), stained with ethidium bromide and photographed under UV light, and their sizes (ca. 1466 bp) were compared with molecular weight markers (100-bp DNA Ladder; Invitrogen Corp., Carlsbad, CA). PCR products were purified using the Illustra GFX PCR DNA and Gel Band Purification kit (GE Healthcare UK Ltd., Buckinghamshire, UK), followed by sequencing at Hokkaido System Science Co. Ltd. (Sapporo, Japan), using the BigDye Terminator Cycle Sequencing kit (Applied Biosystems Japan Ltd., Tokyo, Japan) and an automated DNA sequencer (PRISM-3100, Applied Biosystems Japan Ltd.). Primer 1492 R was used for sequencing (at least 700 bp), and partial 16S rRNA gene sequences were then compared with 16S rRNA gene sequences from the GenBank database using the BLAST search program through the National Center for Biotechnology Information website. Bacterial species were determined by percent sequence similarity ($> 97\%$).

2.6. Determination of bacterial acidogenicity

In order to determine their acidogenicity, bacterial isolates obtained above were cultured on Fastidious Anaerobe Agar (FAA, Lab M, Bury, UK) plates, containing 0.004% bromocresol purple²² in the anaerobic glove box at 37 °C for 2 d. After incubation, bacteria forming yellow ($< \text{pH } 5.2$) zones around their colonies on the plates were termed acidogenic bacteria. On the other hand, those forming purple ($> \text{pH } 6.8$) zones on the plates were termed non-acidogenic bacteria. Furthermore, the acidogenicity of bacterial isolates was confirmed by collecting 1 loopful of each bacterial colony from the CDC anaerobe blood agar plate, followed by culture in 1.0 mL of 1.7% tryptone broth, containing 1.0% glucose in the anaerobic glove box at 37 °C for 2 d. After incubation, pH values of the culture suspensions were measured with a pH meter (pH BOY, Shindengen Electric Manufacturing Co., Ltd., Tokyo, Japan), and were confirmed to be $< \text{pH } 5.2$.

2.7. Data analysis

Tukey's tests were used to analyze differences between data, with P values < 0.05 being considered to be statistically significant.

3. Results

Bacterial growth (log CFU/mg) in plaque samples from molars with orthodontic bands, brackets, and without appliances was 6.6 ± 6.5 , 6.9 ± 7.1 , and 7.4 ± 7.6 , respectively (Table 1). The proportions of facultative anaerobes from molars with orthodontic bands, brackets, and without appliances were 83.3%, 78.3%, and 83.7%, respectively (Table 2). Among the facultative anaerobes, the proportion of *Streptococcus* (44.4%) was higher than that of *Actinomyces* (17.6%) in samples from molars with orthodontic bands ($P < 0.01$). In contrast, *Actinomyces* (43.5 and 40.0%) and *Streptococcus* (23.5 and 34.7%) were more predominant in plaque from molars with orthodontic brackets and without appliances, respectively. Furthermore, the facultative anaerobe, *Neisseria*, was predominant in the plaque microflora from molars with orthodontic bands (Table 2).

Among the obligate anaerobes, *Veillonella* was predominant on molars with orthodontic brackets (8.7%) and without appliances (10.0%), but was scarcely detected on molars with orthodontic bands (1.9%) (Table 2). *Eubacterium* and *Porphyromonas* were specifically detected on molars with orthodontic brackets, but not on those with orthodontic bands and without appliances.

The proportions of acidogenic bacteria were highest on the molars without appliances (81.6%), followed by those with orthodontic bands (74.5%) and brackets (71.3%) (Table 2). The final pH values for culture media containing acidogenic bacteria from molars with orthodontic bands, brackets, and without appliances were 4.5 ± 0.2 , 4.3 ± 0.2 , and 4.3 ± 0.1 , respectively. Most of the acidogenic bacteria were facultative anaerobes, such as *Streptococcus* and *Actinomyces*. The total proportions of *Streptococcus* and *Actinomyces* among acidogenic bacteria were 83.8% (67 of 80 strains), 93.9% (77 of 82), and 91.6% (142 of 155) in samples from molars with orthodontic bands, brackets, and without appliances, respectively. The other acidogenic facultative anaerobes were *Gemella*, *Haemophilus*, and *Actinobacillus*. In contrast, acidogenic obligate anaerobes were *Atopobium* and *Prevotella* (Table 2). There were no particular relationships between sampling period (e.g., before fixation or after removal of orthodontic appliances) and microbial composition.

4. Discussion

Bacterial growth on molars with orthodontic bands was slightly lower than on those with brackets and without appliances (Table 1). Ionomeric materials, particularly glass ionomer cement, have shown antimicrobial properties, which have been attributed to their ability to release fluoride^{23,24}. Furthermore, there are other elements that are simultaneously released from glass ionomer cement²⁵, and these elements may have effects on plaque microflora adjacent to orthodontic bands.

In the present study, the composition of plaque microflora differed depending on surface properties. More specifically, in accordance with a previous study on the composition of plaque microflora on enamel, we found the predominant bacteria in plaque microflora on the surface of molars with orthodontic brackets and without appliances to be *Actinomyces*, *Streptococcus*, and *Veillonella*²⁶. In contrast, in the plaque microflora from molars with orthodontic bands, the proportion of *Streptococcus* was higher, while that of *Actinomyces* and *Veillonella* was lower (Table 2).

In the current study, the proportion of acidogenic bacteria was highest on molars without appliances (81.6%), followed by those with orthodontic bands (74.5%) and brackets (71.3%) (Table 2). Most of the acidogenic bacteria were facultative anaerobes, such as *Streptococcus* (non-mutans streptococci, data not shown) and *Actinomyces*. It is known that non-mutans streptococci and

Table 2

Bacterial composition of plaque biofilm on molars with orthodontic bands, brackets and without appliances.

Subjects	Orthodontic bands								Orthodontic brackets								Without appliances											
	1	1	3	5	5	6	6	Total	2	2	2	3	4	4	4	Total	1	2	2	2	3	3	3	4	4	5	6	Total
Sampled tooth (first molar) ^a	UR	UL	UL	UR	UL	UR	UL		UR	UL	LR	UL	UR	UL	LR		LR	UR	UL	LR	UR	UL	LL	LR	LL	LR	LR	
Total isolates	16	5	17	17	21	17	15	108 (100) ^b	4	18	21	13	21	23	15	115 (100)	11	11	17	21	17	11	18	24	21	24	15	190 (100)
Total of acidogenic bacterial isolates ^c	4	2	16	15	14	15	14	80 (74.5)	2	8	12	12	18	19	11	82 (71.3)	8	11	17	20	10	7	7	19	20	23	13	155 (81.6)
Facultative anaerobes	15	5	11	16	14	15	14	90 (83.3)	3	11	15	12	17	18	14	90 (78.3)	9	11	17	20	11	8	8	19	19	24	13	159 (83.7)
<i>Streptococcus</i> ^c	2	2	10	6	10	10	8	48 (44.4) ^d	2	2	9	6	4	1	3	27 (23.5)	4	10	11	4	7	2	2	5	6	7	8	66 (34.7)
<i>Actinomyces</i> ^c			1	7	1	4	6	19 (17.6) ^e		4	3	6	13	17	7	50 (43.5)			4	16	3	4	5	13	12	16	3	76 (40.0)
<i>Neisseria</i>	9	3						12 (11.1)			1					1 (0.9)							1					1 (0.5)
<i>Gemella</i> ^c	1			2	3	1		7 (6.5)		1					1	2 (1.7)	1	1	2					1	1	2		8 (4.2)
<i>Capnocytophaga</i>	1			1				2 (1.9)	1	3	1				1	6 (5.2)	1				1			1				3 (1.6)
<i>Haemophilus</i> ^c	1							1 (0.9)								0	2											2 (1.1)
<i>Rothia</i>	1							1 (0.9)								0						1						1 (0.5)
<i>Actinobacillus</i> ^c								0								0	1											1 (0.5)
<i>Campylobacter</i>								0		1	1					2 (1.7)												0
<i>Kingella</i>								0							2	2 (1.7)												0
Obligate anaerobes	1	0	6	1	7	2	1	18 (16.7)	1	7	6	1	4	5	1	25 (21.7)	2	0	0	1	6	3	10	5	2	0	2	31 (16.3)
<i>Atopobium</i> ^c			5					5 (4.6)							1	1 (0.9)												0
<i>Peptostreptococcus</i>				1	3		1	5 (4.6)							2	3 (2.6)												0
<i>Slackia</i>					3			3 (2.8)								0												0
<i>Veillonella</i>						1		2 (1.9)	1	2	5	1	1			10 (8.7)	1			3			10	2	1		2	19 (10.0)
<i>Cardiobacterium</i>						1		1 (0.9)								0			1									1 (0.5)
<i>Leptotrichia</i>	1							1 (0.9)							1	1 (0.9)	1			1								2 (1.1)
<i>Olsenella</i>			1					1 (0.9)								0												0
<i>Eubacterium</i>								0					2	1		3 (2.6)												0
<i>Fusobacterium</i>								0		1						1 (0.9)						1		1				2 (1.1)
<i>Porphyromonas</i>								0		2	1					3 (2.6)												0
<i>Prevotella</i> ^c								0		1			1			2 (1.7)								1	1			2 (1.1)
<i>Selenomonas</i>								0		1						1 (0.9)				2	2		1					5 (2.6)

^a UR, upper right; UL, upper left; LR, lower right; LL, lower left.^b Percentages are given in parentheses.^c After two-day incubation, bacteria forming yellow (< pH 5.2) zones around their colonies on fastidious anaerobe agar plates containing 0.004% bromocresol purple were termed acidogenic bacteria.^d Significantly different ($P < 0.01$) between e.^e Significantly different ($P < 0.01$) between d.

Actinomyces are the initial colonizers in plaque biofilm formation²⁷. These findings suggest that the acidogenic potential of plaque microflora upon fixation of orthodontic brackets and/or bands is unexpectedly lower than that of teeth without appliances. Rather, periodontitis-associated bacteria may be more likely to colonize the former plaque microflora (see below).

In the plaque microflora from molars with orthodontic bands, the facultative anaerobe, *Neisseria*, was predominant (Table 2). In contrast, in microflora from molars with brackets or without any orthodontic appliances, the obligate anaerobe, *Veillonella*, known as a late colonizer in dental plaque formation, was predominant. It has been reported that *Neisseria* species colonize and constitute 1.3% to 15.2% of the early plaque biofilm on enamel²⁸. These findings suggest that the partial pressure of oxygen may be higher in plaque that forms close to metal, possibly due to the plaque structure being less mature²⁹. In turn, this would result in the low proportions of obligate anaerobes found in plaque on molars with orthodontic bands (Table 2), similar to that reported in shallow periodontal pockets³⁰.

Eubacterium, *Porphyromonas*, *Fusobacterium*, and *Prevotella* were detected in the plaque on molars with orthodontic brackets, as well as from those without appliances (Table 2), but were not detected on molars with orthodontic bands. It has been reported that these obligate anaerobes are one of the components of mature plaque microflora^{2,31,32}, and that they are associated with periodontitis^{33,34}. In addition, *Prevotella intermedia* and *Tannerella forsythia* have frequently been reported to be present in subgingival plaque after fixation of orthodontic brackets^{13,35}. Furthermore, the CFU ratio of anaerobic bacteria has been reported to increase after fixation of orthodontic appliances^{36–39}. These findings suggest that plaque microflora on molars with orthodontic brackets is similar to that of periodontitis-associated microflora. Due to the complex structure of brackets, it is possible that oxygen scarcely infiltrates into dental plaque on the surfaces of teeth with orthodontic brackets. Thus, establishment of anaerobic environments in this way can promote obligate anaerobes, including periodontitis-associated bacteria, such as *Eubacterium*, *Porphyromonas*, *Fusobacterium*, and *Prevotella*, to colonize and grow^{3,30,40}.

We found no particular relationship between sampling sites (e.g., upper and lower first molars) and microbial composition. Furthermore, since relatively few plaque samples were examined in this study, correlations between age and ecologic distribution of oral microbiota remain uncertain, and further large-scale studies are necessary.

5. Conclusions

The findings of the present study indicate that there are differences in the microbial composition and acidogenic potential of supragingival plaque on first molars with orthodontic bands, brackets, and without appliances. Our data further suggest that supragingival plaque on teeth with brackets carries higher levels of periodontitis-associated bacteria but less caries-associated bacteria.

Conflict of interest

No potential conflicts of interest are disclosed.

Acknowledgments

This study was supported in part by Grants-in-Aid for Scientific Research (22390399, 23592791) from the Japan Society for the Promotion of Science.

References

- [1] Kolenbrander PE, Palmer Jr RJ, Periasamy S, Jakubovics NS. Oral multispecies biofilm development and the key role of cell-cell distance. *Nat Rev Microbiol* 2010;8:471–80.
- [2] Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, Sahasrabudie A, Dewhirst FE. Bacterial diversity in human subgingival plaque. *J Bacteriol* 2001;183:3770–83.
- [3] Takahashi N. Microbial ecosystem in the oral cavity: Metabolic diversity in an ecological niche and its relationship with oral diseases. *Int Cong Ser* 2005;1284:103–12.
- [4] O'Reilly MM, Featherstone JDB. Demineralization and remineralization around orthodontic appliances: an in vivo study. *Am J Orthod Dentofac Orthop* 1987;92:33–40.
- [5] Sanpei S, Endo T, Shimooka S. Caries risk factors in children under treatment with sectional brackets. *Angle Orthod* 2010;80:509–14.
- [6] Pramod S, Kailasam V, Padmanabhan S, Chitharanjan AB. Presence of cariogenic streptococci on various bracket materials detected by polymerase chain reaction. *Aust Orthod J* 2011;27:46–51.
- [7] Nelson-Filho P, Olmedo LY, Andruccioli MC, Saraiva Mda C, Matsumoto MA, de Queiroz AM, da Silva RA, da Silva LA. Use of the checkerboard DNA–DNA hybridisation technique for in vivo detection of cariogenic microorganisms on metallic brackets, with or without use of an antimicrobial agent. *J Dent* 2011;39:513–7.
- [8] Türkkanhraman H, Sayin MÖ, Bozkurt FY, Yetkin Z, Kaya S, Önal S. Archwire ligation techniques, microbial colonization, and periodontal status in orthodontically treated patients. *Angle Orthod* 2005;75:231–6.
- [9] Ai H, Lu HF, Liang HY, Wu J, Li RL, Liu GP, Xi Y. Influences of bracket bonding on mutans streptococcus in plaque detected by real time fluorescence-quantitative polymerase chain reaction. *Chin Med J* 2005;118:2005–10.
- [10] Anhoury P, Nathanson D, Hughes CV, Socransky S, Feres M, Chou LL. Microbial profile on metallic and ceramic bracket materials. *Angle Orthod* 2002;72:338–43.
- [11] Kim SH, Choi DS, Jang I, Cha BK, Jost-Brinkmann PG, Song JS. Microbiologic changes in subgingival plaque before and during the early period of orthodontic treatment. *Angle Orthod* 2012;82:254–60.
- [12] Demling A, Demling C, Schweska-Polly R, Stiesch M, Heuer W. Short-term influence of lingual orthodontic therapy on microbial parameters and periodontal status. A preliminary study. *Angle Orthod* 2010;80:480–4.
- [13] Choi DS, Cha BK, Jost-Brinkmann PG, Lee SY, Chang BS, Jang I, Song JS. Microbiologic changes in subgingival plaque after removal of fixed orthodontic appliances. *Angle Orthod* 2009;79:1149–55.
- [14] Thornberg MJ, Riolo CS, Bayirli B, Riolo ML, Van Tubergen EA, Kulbersh R. Periodontal pathogen levels in adolescents before, during, and after fixed orthodontic appliance therapy. *Am J Orthod Dentofac Orthop* 2009;135:95–8.
- [15] Lee SM, Yoo SY, Kim HS, Kim KW, Yoon YJ, Lim SH, Shin HY, Kook JK. Prevalence of putative periodontopathogens in subgingival dental plaques from gingivitis lesions in Korean orthodontic patients. *J Microbiol* 2005;43:260–5.
- [16] Sallum EJ, Nouer DF, Klein MI, Gonçalves RB, Machion L, Wilson Sallum A, Sallum EA. Clinical and microbiologic changes after removal of orthodontic appliances. *Am J Orthod Dentofac Orthop* 2004;126:363–6.
- [17] Stecksén-Blicks C, Holgersson PL, Olsson M, Bylund B, Sjöström I, Sköld-Larsson K, Kalfas S, Twetman S. Effect of xylitol on mutans streptococci and lactic acid formation in saliva and plaque from adolescents and young adults with fixed orthodontic appliances. *Eur J Oral Sci* 2004;112:244–8.
- [18] Sengun A, Sari Z, Ramoglu SI, Malkoç S, Duran I. Evaluation of the dental plaque pH recovery effect of a xylitol lozenge on patients with fixed orthodontic appliances. *Angle Orthod* 2004;74:240–4.
- [19] Sköld-Larsson K, Borgström MK, Twetman S. Effect of an antibacterial varnish on lactic acid production in plaque adjacent to fixed orthodontic appliances. *Clin Oral Invest* 2001;5:118–21.
- [20] Lane DJ. 16 S/23 S rRNA sequencing. In: Stackebrandt E, Goodfellow M, editors. *Nucleic acid techniques in bacterial systematics*. Chichester, UK: John Wiley & Sons; 1991. p. 115–75.
- [21] Marchesi JR, Sato T, Weightman AJ, Martin TA, Fry JC, Hiom SJ, Dymock D, Wade WG. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16 S rRNA. *Appl Environ Microbiol* 1998;64:795–9.
- [22] Smith CA, Venning M, Verran J. Use of novel culture media to indicate alkaligenic properties of dental plaque. *Microb Ecol Health Dis* 2004;16:44–50.
- [23] Forss H, Jokinen J, Spets-Happonen S, Seppä L, Luoma H. Fluoride and mutans streptococci in plaque grown on glass ionomer and composite. *Caries Res* 1991;25:454–8.
- [24] Benelli EM, Serra MC, Rodrigues Jr. AL, Curly JA. In situ anticariogenic potential of glass ionomer cement. *Caries Res* 1993;27:280–4.
- [25] Nakajo K, Imazato S, Takahashi Y, Kiba W, Ebisu S, Takahashi N. Fluoride released from glass-ionomer cement is responsible to inhibit the acid production of caries-related oral streptococci. *Dent Mater* 2009;25:703–8.
- [26] Hoshino E, Sato M, Sasano T, Kota K. Characterization of bacterial deposits formed in vivo on hydrogen-ion-sensitive field-effect transistor electrodes and enamel surfaces. *Jpn J Oral Biol* 1989;31:102–6.
- [27] Kolenbrander PE, Andersen RN, Clemans DL, Whittaker CJ, Klier CM. Potential role of functionally similar coaggregation mediators in bacterial succession.

- In: Newman HN, Wilson M, editors. *Dental Plaque Revisited*. Cardiff, UK: BioLine; 1999. p. 171–86.
- [28] Diaz PI, Chalmers NI, Rickard AH, Kong C, Milburn CL, Palmer Jr. RJ, Kolenbrander PE. Molecular characterization of subject-specific oral microflora during initial colonization of enamel. *Appl Environ Microbiol* 2006;72:2837–48.
- [29] Quirynen M, Listgarten MA. The distribution of bacterial morphotypes around natural teeth and titanium implants ad modum Brånemark. *Clin Oral Implants Res* 1990;1:8–12.
- [30] Loesche WJ, Gusberti F, Mettraux G, Higgins T, Syed S. Relationship between oxygen tension and subgingival bacterial flora in untreated human periodontal pockets. *Infect Immun* 1983;42:659–67.
- [31] Mayanagi G, Sato T, Shimauchi H, Takahashi N. Detection frequency of periodontitis-associated bacteria by polymerase chain reaction in subgingival and supragingival plaque of periodontitis and healthy subjects. *Oral Microbiol Immunol* 2004;19:379–85.
- [32] Uematsu H, Hoshino E. Predominant obligate anaerobes in human periodontal pockets. *J Periodontol Res* 1992;27:15–9.
- [33] Moore WEC, Moore LVH. The bacteria of periodontal diseases. *Periodontol* 2000 1994;5:66–77.
- [34] Wade WG, Munson MA, de Lillo A, Weightman AJ. Specificity of the oral microflora in dental caries, endodontics infections and periodontitis. *Int Cong Ser* 2005;1284:150–7.
- [35] Ristic M, Vlahovic Svabic M, Sasic M, Zelic O. Effects of fixed orthodontic appliances on subgingival microflora. *Int J Dent Hyg* 2008;6:129–36.
- [36] van Gastel J, Teughels W, Quirynen M, Struyf S, van Damme J, Coucke W, Carels C. Longitudinal changes in gingival crevicular fluid after placement of fixed orthodontic appliances. *Am J Orthod Dentofac Orthop* 2011;139:735–44.
- [37] van Gastel J, Quirynen M, Teughels W, Coucke W, Carels C. Longitudinal changes in microbiology and clinical periodontal parameters after removal of fixed orthodontic appliances. *Eur J Orthod* 2011;33:15–21.
- [38] van Gastel J, Quirynen M, Teughels W, Coucke W, Carels C. Longitudinal changes in microbiology and clinical periodontal variables after placement of fixed orthodontic appliances. *J Periodontol* 2008;79:2078–86.
- [39] van Gastel J, Quirynen M, Teughels W, Coucke W, Carels C. Influence of bracket design on microbial and periodontal parameters in vivo. *J Clin Periodontol* 2007;34:423–31.
- [40] Takahashi N, Saito K, Schachtele CF, Yamada T. Acid tolerance and acid-neutralizing activity of *Porphyromonas gingivalis*, *Prevotella intermedia* and *Fusobacterium nucleatum*. *Oral Microbiol Immunol* 1997;12:323–8.